Trapping state restoration in the randomly-driven Jaynes-Cummings model by conditional measurements

Mauro Fortunato*[(a)], Gershon Kurizki[(b)], and Wolfgang P. Schleich[(a)]
(a) Abteilung für Quantenphysik, Universität Ulm, Albert-Einstein-Allee 11, D-89069 Ulm, Germany
(b) Department of Chemical Physics, The Weizmann Institute of Science, Rehovot 76100, Israel
(February 9, 2008)

arXiv:quant-ph/9808049v2 16 Sep 1998

We propose a scheme which can effectively restore fixed points in the quantum dynamics of repeated Jaynes-Cummings interactions followed by atomic state measurements, when the interaction times fluctuate randomly. It is based on selection of superposed atomic states whose phase correlations tend to suppress the phase fluctuations of each separate state. One suggested realization involves the convergence of the cavity field distribution to a single Fock state by conditional measurements performed on two-level atoms with fluctuating velocities after they cross the cavity. Another realization involves a trapped ion whose internal-motional state coupling fluctuates randomly. Its motional state is made to converge to a Fock state by conditional measurements of the internal state of the ion.

PACS numbers: 42.50.-p,42.50.Dv,32.80.-t,05.45.+b

It is our purpose here to propose a method of controlling the evolution of a randomly driven system by means of quantum measurements. The basic idea is that in an ensemble of randomly phased systems it is possible to select (project out), by appropriate quantum measurements, a *superposition* of two (or more) sub-ensembles with a *fixed phase difference*. As a result, the phase fluctuations of the selected superposed sub-ensembles can be suppressed by their interference as compared to the separate fluctuations of each sub-ensemble.

The manipulation of the system evolution by first unitarily entangling it with another system (a "meter"), and then post-selecting a certain superposition of eigenstates of the meter, is known as the conditional measurement (CM) approach [1]. This approach has been developed for the purpose of controlling the quantum state of the cavity field by its interactions with atoms, followed by post-selection of chosen atomic-eigenstate superpositions [1]. The price one has to pay for guiding the evolution by CMs is that the success probability of every CM, which is measured by the *fraction* of post-selected states among all possible measurement outcomes, is by necessity less than unity. Hence, one must perform as many trials as implied by the success probability until the required CM is accomplished (see below).

Although the outlined proposal for controlling a randomly-driven system by CMs is quite general, we shall apply it here specifically to the fundamentally simple Jaynes-Cummings model (JCM) [1,2] and its classical counterparts. This model describes the interaction of a two-level dipole transition, $|e\rangle \longleftrightarrow |g\rangle$ with a resonant quantized field mode, via the secular (rotating-wave) interaction Hamiltonian

$$H_{\rm int} = g(|e\rangle\langle g|a + {\rm h.c.}) \ , \tag{1}$$

g being the field-dipole coupling strength and a the field annihilation operator. This interaction can be viewed as the resonant coupling of two quantized pendula. The merits of this model are that it is exactly solvable (in the secular, or rotating-wave approximation) and is experimentally realizable for atoms crossing a high-Q cavity [1–5] or for trapped ions interacting with standing laser waves [6,7].

We first discuss the cavity realization of this model. In the interaction picture, the solution of the JCM for an atom initially in the excited state $|e\rangle$ and the field in a superposition of many Fock (photon-number) states, $|\psi_{\rm F}^{(i)}\rangle = \sum_n c_n^{(i)}|n\rangle$, after an interaction time τ, is given by

$$\begin{aligned} |\psi_{\rm AF}\rangle &= \hat{U}(\tau)|e\rangle|\psi_{\rm F}^{(i)}\rangle \\ &= \sum_n c_n^{(i)}\left[\cos\theta_n|e\rangle|n\rangle - i\sin\theta_n|g\rangle|n+1\rangle\right] \ , \end{aligned} \tag{2}$$

where $\hat{U}(\tau)$ is the JCM evolution operator and $\theta_n = g\tau\sqrt{n+1}$. This solution describes a superposition of sinusoidal energy exchanges between the "pendula", corresponding to the n-dependent amplitudes of stimulated photon emission and reabsorption by the atom as a function of its flight time τ through the cavity. This description neglects dissipation, which is justifiable when the lifetimes of state $|e\rangle$ and of state $|\psi_{\rm F}^{(i)}\rangle$ in the cavity are sufficiently long, as discussed below. The essential feature of Eq. (2) is that it describes the entanglement of the field and atom states via $\hat{U}(\tau)$.

Our study will focus on the effect of fluctuations in the atomic velocity, and therefore in τ, on the evolution of the field state, which is driven by successive interactions with atoms via Eq. (2), followed by measurements performed on the atoms after exiting the cavity. It has been noted [2,3,8] that such evolution can cause the convergence of an initial distribution of Fock states to particular

Fock states which are the "trapping states" of the unitary transformation (2), satisfying the condition $\theta_n = q\pi$, q being an integer. The originally proposed convergence procedure relied on non-selective measurements (NSM) of the atoms, *i.e.*, on ignoring the final state of each atom exiting the cavity, and merely recording its presence [8]. If the Fock state distribution after the passage of $k-1$ atoms through the cavity is $\rho^{(k-1)}_{\mathrm{F},nn} = |c^{(k-1)}_n|^2$, then the entanglement with the kth atom according to Eq. (2) followed by tracing over the final atomic states results in the transformation

$$|c_n^{(k-1)}|^2 \rightarrow |c_n^{(k-1)}|^2 \cos^2\theta_n^{(k)} + |c_{n-1}^{(k-1)}|^2 \sin^2\theta_{n-1}^{(k)} \,. \quad (3)$$

This transformation implies that each Fock state $|n\rangle$ loses part of its population to $|n+1\rangle$ by stimulated emission, *unless* $\sin^2\theta^{(k)}_{n-1} = 0$, in which case it becomes an attractor or "trapping" state, whose population grows at the expense of Fock states with *lower* photon numbers. Thus, a Fock-state distribution localized *below* a trapping state $|n_t\rangle$ should ideally converge to the photon-number distribution δ_{n,n_t}, if all atomic flight times τ_k are equal and exactly satisfy the above "trapping" (attractor) condition. However, such an attractor is merely a *marginally stable* fixed point, *i.e.*, it is stable when approached from below ($n < n_t$), but unstable for $n > n_t$. This stability is very fragile, and is destroyed even by very small fluctuations in the interaction times around the trapping condition. This is shown in Fig. 1(a) and (b), where we plot as a function of the number of atoms injected the mean value and the rms value of the photon number, in the case (a) of fixed interaction times and (b) of tiny (1%) time fluctuations.

These stability properties of the trapping Fock states go over smoothly into their classical counterparts, in the limit wherein the initial cavity field is nearly classical, allowing the replacement of the JCM evolution [Eq. (2)] by that of the parametrically driven classical pendulum [8]

$$\frac{\partial\theta}{\partial(g\tau)} = \epsilon, \qquad \frac{\partial\epsilon}{\partial(g\tau)} = \sin\theta \,, \quad (4)$$

where θ is the atomic polarization (the pendulum tipping angle), ϵ is the dimensionless cavity field, and the atoms are injected into the cavity in the excited (inverted) state. The return map $\epsilon_{k+1} = f(\epsilon_k)$ corresponding to Eq. (4) allows us to plot the evolution of the field-energy distribution for fluctuating τ_k. Under the assumption of small changes in the classical field due to the interaction with one atom, this return map can be approximated by [8]

$$\epsilon_{k+1} \approx \epsilon_k + \frac{2}{\epsilon_k}\sin^2\left(\frac{\epsilon_k g\tau_k}{2}\right) \,. \quad (5)$$

The marginal stability of the fixed points, just as in the quantum domain, makes the convergence practically impossible, and leads to *intermittently chaotic* behavior, even for small τ_k fluctuations. This is clearly shown in Fig. 1(c) and (d), where we plot as a function of the number of atoms the classical counterpart $\epsilon^2/4$ of the mean number of photons $\langle n\rangle$, in the case of fixed interaction times, and of tiny (1%) fluctuations, respectively.



FIG. 1. Sensitivity of the NSM scheme (both quantum and classical) to time-fluctuations. Top: behavior of $\langle n\rangle$ and Δn as a function of the number of injected atoms in the quantum case for (a) fixed interaction times ($|n_t\rangle = |138\rangle$) and (b) tiny (1%) fluctuations around the trapping value; the initial coherent state is $|\alpha\rangle = |3\rangle$. Bottom: evolution of $\epsilon^2/4$ (the classical counterpart of $\langle n\rangle$) versus number of atoms in the classical case for (c) fixed interaction times ($g\tau = 2\pi/\sqrt{199}$) and (d) tiny (1%) fluctuations around this value; initially, $\epsilon^2/4 = 9$.

Are these fixed points irretrievable in the presence of τ-fluctuations? In order to examine this question, we consider a different kind of field-state transformations, resulting from atomic CMs, since such transformations can generally better guide the field evolution towards particular states [1]. Schematically, k CMs are performed on the cavity field state if each of the k atoms that have crossed the cavity is successfully detected in the desired final state $|\varphi^{(f)}_k\rangle$. The resulting field-state transformation $|\psi_{k-1}\rangle \rightarrow |\psi_k\rangle$ is effected by the kth CM according to

$$|\psi_k\rangle = P_k^{-1/2}\left[\langle\varphi_k^{(f)}|\hat{U}(\tau_k)\left(|\varphi_k^{(i)}\rangle|\psi_{k-1}\rangle\right)\right] \,, \quad (6)$$

where $\hat{U}(\tau_k)$ entangles the initial atomic state $|\varphi^{(i)}_k\rangle$ with the field state $|\psi_{k-1}\rangle$, and the CM success probability P_k is the norm of the expression in square brackets. The simplest choice of final atomic states $|\varphi^{(f)}_k\rangle$ onto which the entangled state is projected appears to be one of the energy states, $|e\rangle$ or $|g\rangle$. For atoms injected initially in $|\varphi^{(i)}_k\rangle = |e\rangle$, CMs consisting in post-selection of $|\varphi^{(f)}_k\rangle = |e\rangle$ or $|g\rangle$ can be referred to as the "elastic" and "inelastic" selected sub-ensembles, respectively [1,8]. Does the selection of one such sub-ensemble improve the



FIG. 2. Evolution of $\langle n\rangle$ and Δn versus k (number of atoms) in the scheme of elastic CMs for (a) small spread in the interaction times ($\Delta\tau = \Delta\tau_c/10$), and (b) large spread ($\Delta\tau = \Delta\tau_c$). The initial coherent state is $|\alpha\rangle = |3\rangle$. The random fluctuations are around the value $\tau_k = \pi/g\sqrt{21}$, and therefore the trapping state (for fixed interaction time) is $|n_t\rangle = |20\rangle$.

convergence of a distribution of Fock states to a trapping state $|n_t\rangle$ when τ_k fluctuate? As seen in Fig. 2, it does, as long as the spread of τ_k is below the critical value $\Delta\tau_c$. This critical value can be estimated as the difference between the trapping ($\theta_{n_t} = \pi$) and the anti-trapping ($\theta_{n_t} = \pi/2$) interaction times. For the "elastic" sub-ensemble $|\varphi_k^{(i)}\rangle = |e\rangle \rightarrow |\varphi_k^{(f)}\rangle = |e\rangle$ this yields $\Delta\tau_c \simeq \pi/(g\sqrt{n_t+1}) - \pi/(2g\sqrt{n_t+1})$. This estimate is fully confirmed by the numerical results in Fig. 2: whereas for τ_k-spreads $\Delta\tau \simeq \Delta\tau_c/10$ convergence is still possible, it fails completely for $\Delta\tau = \Delta\tau_c$, and the system escapes its fixed points because the trapping condition is different for each atom.

These findings can be carried over to the quasi-classical domain: The selection of the "elastic" sub-ensemble ($|e\rangle \rightarrow |e\rangle$) corresponds to singling out atoms that do not change, on average, the cavity field energy by virtue of trapping. This selection should help keep the field energy near a fixed point, either classically or quantum mechanically, but only as long as the fluctuations about the fixed-point (trapping) condition are small.

However, instead of "elastic" or "inelastic" post-selection, it is possible to design CMs which post-select any desired superpositions of the above sub-ensembles [9]. Such post-selections can be experimentally realized [5] by passing the atoms after the cavity through a classical field (Ramsey zone) setup over a time-interval T_k, which rotates, say, the final state as follows

$$|e\rangle \longrightarrow |\varphi_k^{(f)}\rangle = \alpha_k^{(f)}|e\rangle + \beta_k^{(f)}|g\rangle \ , \tag{7a}$$

$$\alpha_k^{(f)} \ = \ \cos(\Omega T_k/2) \ , \ \ \beta_k^{(f)} = \sin(\Omega T_k/2) e^{i\phi_f} \ , \tag{7b}$$

Ω being the Rabi frequency of the classical field and ϕ_f an *independent* phase shift. The corresponding CM transformation [Eq. (6)] $|\psi_{k-1}\rangle = \sum_n c_n^{(k-1)}|n\rangle \rightarrow |\psi_k\rangle = \sum_n c_n^{(k)}|n\rangle$ can be written as

$$c_n^{(k-1)} \rightarrow c_n^{(k)} = P_k^{-1/2}\Big[\alpha_k^{(f)*}\cos\theta_n^{(k)} c_n^{(k-1)} - i\beta_k^{(f)*}\sin\theta_{n-1}^{(k)} c_{n-1}^{(k-1)}\Big] \ . \tag{8}$$



FIG. 3. Stability of a Fock state—even for a large spread in the interaction times—in the post-selection scheme $|e\rangle \rightarrow |e\rangle + |g\rangle$ for $|n_t\rangle = |21\rangle$. Top: case of small fluctuations in the interaction times ($\Delta\tau = \Delta\tau_c/10$): behavior of (a) $\langle n\rangle$ and of (b) Δn versus k (number of atoms). Bottom: the same for large spread in the interaction times ($\Delta\tau = 2\Delta\tau_c$): (c) $\langle n\rangle$ and (d) Δn. In all cases the initial coherent state is $|\alpha\rangle = |\sqrt{21}\rangle$.

If we neglect the differences between $\theta_n^{(k)}$ and $\theta_{n-1}^{(k)}$ or between $c_n^{(k-1)}$ and $c_{n-1}^{(k-1)}$, consistently with the large n limit, and use the explicit forms of $\alpha_k^{(f)}$ and $\beta_k^{(f)}$ for a Ramsey-zone field, choosing $\phi_f = -\pi/2$, then Eq. (8) becomes

$$c_n^{(k)} \simeq P_k^{-1/2}\cos\left(\Omega T_k/2 - g\tau_k\sqrt{n+1}\right) c_n^{(k-1)} \ . \tag{9}$$

Since the atoms, whose velocities fluctuate, first cross the cavity and then the classical field zone, T_k and τ_k in Eq. (9) are correlated, although random. This correlation, which results from the *interference* of the post-selected sub-ensembles, yields a *strong suppression* of the cosine fluctuations in Eq. (9). The consequences of this suppression are shown in Fig. 3 where we plot (for the scheme $|e\rangle \rightarrow |e\rangle + |g\rangle$) $\langle n\rangle$ and Δn for small and large spreads in the interaction times: Despite the large fluctuations, the convergence towards the desired trapping state is very good.

In Fig. 4 the photon-number distribution after 2000 atoms is plotted for the case of Fig. 3(c) and (d). This confirms that the state produced inside the cavity is indeed the Fock state $|n\rangle = |21\rangle$: all the $P(n)$ vanish apart from one, $P(21) = 1$.

The above analysis demonstrates that the superposed-subensembles CM allows the restoration of field trapping states in simple dynamical evolution, as effected by the resonant JCM. The price one has to pay for this effect is given by the success probability of the sequence of CMs, which amounts to post-selecting the desired sub-ensemble and discarding all other sequences. For fixed interaction times the success probabilities $\mathcal{P}_N$ of the whole sequence



FIG. 4. Final photon-number distribution after $N = 2000$ atoms in the case of $|e\rangle \to |e\rangle + |g\rangle$ CM scheme, for large fluctuations in the interaction times.

of N atoms, even though it is monotonically decreasing, converges towards the norm of the overlap between the initial (coherent) state and the trapping Fock state. In the case of fluctuating times, this value is significantly smaller. Still, for the parameters of Fig. 3(c) and (d) and for $N = 110$, which suffices for the convergence, we have $\mathcal{P}_N \gtrsim 10^{-3}$.

The random distribution for the interaction times τ_k we have used in our calculation was uniform within $\Delta\tau$. We have also made numerical runs using Gaussian distributions, with the same rms values, obtaining similar results. The only difference, is that in the latter case it can happen—even though with small probability—that one atom crosses the cavity with an interaction time which is far at the distribution tails. This may cause the field evolution to escape the fixed point. Different initial photon-number distributions lead to similar results.

We notice that our results may be of value in the task of state preparation [1–5,8,9]. In particular, they may help in the generation of Fock states, since the present scheme is quite immune even to large fluctuations in the field-atom interaction times.

The most stringent requirement on the realization of our scheme is that the damping time $\tau_{\rm cav}$ of the cavity be much greater than the duration $\tau_{\rm seq}$ of the atomic sequence multiplied by n_t. A simple estimate [1] shows that for $N \simeq 10^3$ atoms with thermal velocity ($v_{\rm at} \sim 10^3$ m/s) crossing a cavity 0.5 cm long, we need a cavity quality factor $Q \gg 10^{10}$, which is currently achievable in superconducting microwave cavities [5].

The above scheme can be fully implemented for a trapped cold ion interacting with traveling- and standing-wave laser fields [6]. The JCM evolution Eq. (2) then holds for the resonant laser-induced coupling of the internal-state transition $|e\rangle \longleftrightarrow |g\rangle$ with the vibrational state of the ion motion [6]. The transformation of Fock states of the motion $|n\rangle$ [Eq. 3] is then determined by the phases $\theta_n^{(k)} = g\tau_k\sqrt{n+1}$, where τ_k is the duration of the k-th standing-wave laser pulse. The rotation of $|e\rangle$ and $|g\rangle$ into the superpositions given by Eqs. (7) is achieved by means of a traveling-wave laser pulse, whose duration T_k is *correlated* to that of the standing-wave pulse τ_k. Successful CMs, corresponding to Eq. (9), can be performed by detecting the fluorescence emitted by the rotated excited state (7a). These CMs can suppress the adverse effect of laser-pulse fluctuations on the convergence of the motional state to the trapping state $|n\rangle$, just like in the cavity realization. The experimental difficulties of the ion trap realization [7] appear to be much less than those of the cavity realization.

To sum up, in the present scheme, the CMs force the field to evolve towards the trapping state, in spite of the large fluctuations in the interaction times, by using the correlations existing between fluctuating interaction times, and measuring-device ("meter") times. Hence, our scheme can be considered as a novel method of restoring fixed points in the quantum dynamics of a system which is intermittently chaotic, even under large, random perturbations [10–12].

ACKNOWLEDGMENTS

We acknowledge helpful and stimulating discussions with P. Stifter, and the support of the German-Israeli Foundation. M.F. thanks the European Community (Human Capital and Mobility programme) for support.

---

* E-mail: for@physik.uni-ulm.de

[1] The conditional measurement approach has been suggested by B. Sherman and G. Kurizki, Phys. Rev. A **45**, R7674 (1992) and further developed by B. M. Garraway, B. Sherman, H. Moya-Cessa, P. L. Knight, and G. Kurizki, *ibid.* **49**, 535 (1994). One remarkable enrichment of this approach is "quantum state engineering" introduced by K. Vogel, V. M. Akulin, and W. P. Schleich, Phys. Rev. Lett. **71**, 1816 (1993).

[2] E. T. Jaynes and F. W. Cummings, Proc. IEEE **51**, 89 (1963).

[3] J. Krause, M. O. Scully, and H. Walther, Phys. Rev. A **36**, 4547 (1987); P. Meystre, G. Rempe, and H. Walther, Opt. Lett. **13**, 1078 (1988); F. W. Cummings and A. K. Rajagopal, Phys. Rev. A **39**, 3414 (1989); P. Meystre in *Progress in Optics XXX*, E. Wolf (ed.), Elsevier (1992).

[4] See, for example, G. J. Milburn and D. F. Walls, *Quantum Optics* (Springer Verlag, Berlin, 1994), or Ref. [3].

[5] G. Rempe, F. Schmidt-Kaler, and H. Walther, Phys. Rev. Lett. **64**, 2783 (1990); D. Meschede, Phys. Rep. **211**, 201 (1992).

[6] J. I. Cirac, A. S. Parkins, R. Blatt, and P. Zoller, Phys. Rev. Lett. **70**, 556 (1993); J. I. Cirac, R. Blatt,

A. S. Parkins, and P. Zoller, Phys. Rev. Lett. **70**, 762 (1993)

[7] D. M. Meekhof, C. Monroe, B. E. King, W. M. Itano, and D. J. Wineland, Phys. Rev. Lett. **76**, 1796 (1996); C. Monroe, D. M. Meekhof, B. E. King, and D. J. Wineland, Science **272**, 1131 (1996).

[8] P. Filipowicz, J. Javainen, and P. Meystre, J. Opt. Soc. Am. B **3**, 906 (1986). For the measurements-induced dynamics of a micromaser, see: P. Meystre, Opt. Lett. **12**, 669 (1987); P. Meystre and E.M. Wright, Phys. Rev. A **37**, 2524 (1988).

[9] G. Harel, G. Kurizki, J. K. McIver, and E. Coutsias, Phys. Rev. A **53**, 4534 (1996).

[10] H. G. Schuster, *Deterministic Chaos*, 2nd ed. (VCH, Weinheim, 1989); E. Ott, *Chaos in dynamical systems* (Cambridge University Press, 1993).

[11] K. Pyragas, Phys. Lett. A **170**, 421 (1992); S. Rajasekar and M. Lakshmanan, Physica D **67**, 282 (1993); F. T. Arecchi, G. Basti, S. Boccaletti, and A. L. Perrone, Europhys. Lett. **26**, 327 (1994);

[12] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics* (Springer, Berlin, 1990); P. W. Milonni, M. L. Shih, and J. R. Ackerhalt, *Chaos in Laser-Matter Interactions* (World Scientific, Singapore, 1987); R. Schack and C. M. Caves, Phys. Rev. Lett. **69**, 3413 (1992); *ibid.* **71**, 525 (1993).